

Prevented mortality and greenhouse gas emissions from historical and projected nuclear power

Pushker A. Kharecha, and James E Hansen

Environ. Sci. Technol., **Just Accepted Manuscript** • DOI: 10.1021/es3051197 • Publication Date (Web): 15 Mar 2013

Downloaded from <http://pubs.acs.org> on April 2, 2013

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.



Prevented mortality and greenhouse gas emissions from historical and projected nuclear power

Pushker A. Kharecha^{1,*}, James E. Hansen¹

Affiliations:

¹NASA Goddard Institute for Space Studies and Columbia University Earth Institute, 2880 Broadway, New York, NY 10025 USA

*Corresponding author: pushker@giss.nasa.gov, ph: 212-678-5536, fax: 212-678-5552

Abstract

In the aftermath of the March 2011 accident at Japan's Fukushima Daiichi nuclear power plant, the future contribution of nuclear power to the global energy supply has become somewhat uncertain. Because nuclear power is an abundant, low-carbon source of base-load power, on balance it could make a large contribution to mitigation of global climate change and air pollution. Using historical production data, we calculate that global nuclear power has prevented about 1.84 million air pollution-related deaths and 64 gigatonnes (Gt) CO₂-equivalent greenhouse gas (GHG) emissions that would have resulted from fossil fuel burning. Based on global projection data that take into account the effects of Fukushima, we find that by mid-century, nuclear power could prevent an additional 420,000 to 7.04 million deaths and 80 to 240 GtCO₂-eq emissions due to fossil fuels, depending on which fuel it replaces. By contrast, we assess that large-scale expansion of natural gas use would not mitigate the climate problem and would cause far more deaths than expansion of nuclear power.

Introduction

It has become increasingly clear that impacts of unchecked anthropogenic climate change due to GHG emissions from burning of fossil fuels could be catastrophic for both human society and natural ecosystems (in ref 1, see Figures SPM.2 and 4.4), and that the key time frame for mitigating the climate crisis is the next decade or so^{2,3}. Likewise, during the past decade outdoor air pollution due largely to fossil fuel burning is estimated to cause over 1 million deaths annually worldwide⁴. Nuclear energy (and other carbon-free energy sources) could help to mitigate both of these major problems⁵.

The future of global nuclear power will depend largely on choices made by major energy-using countries in the next decade or so⁶. While most of the highly nuclear-dependent countries have affirmed their plans to continue development of nuclear power after the Fukushima accident, several have announced that they will either temporarily suspend plans for new plants or completely phase out existing plants². Serious questions remain about safety, proliferation, and disposal of radioactive waste, which we have discussed in some detail elsewhere⁷.

Here, we examine the historical and potential future role of nuclear power with respect to prevention of air pollution-related mortality as well as GHG emissions on multiple spatial scales. Previous studies have quantified global-scale avoided GHG emissions due to nuclear power (e.g., refs 5, 8-10), however the issue of avoided human deaths remains largely unexplored. We focus on the world as a whole, OECD Europe, and the five countries with the highest annual CO₂ emissions in the last several years. In order, these top five CO₂ emitters are China, United States,

1 India, Russia, and Japan, accounting for 56% of global emissions from 2009-2011¹¹. To estimate
 2 historically prevented deaths and GHG emissions, we start with data for global annual electricity
 3 generation by energy source from 1971-2009 (Figure 1). We then apply mortality and GHG
 4 emissions factors, defined respectively as deaths and emissions per unit electric energy
 5 generated, for relevant electricity sources (Table 1). For the projection period (2010-2050), we
 6 base our estimates on recent (post-Fukushima) nuclear power trajectories given by the UN
 7 International Atomic Energy Agency (IAEA)⁶.

11 Methods

14 Calculation of prevented mortality and GHG impacts

15 For the historical period (1971-2009), we assume that all nuclear power supply in a given
 16 country and year would instead have been delivered by fossil fuels (specifically coal and natural
 17 gas), given their worldwide dominance and the very small contribution of non-hydro renewables
 18 to world electricity thus far (Figure 1). There are, of course, numerous complications involved in
 19 trying to design such a replacement scenario (e.g., evolving technological and socioeconomic
 20 conditions), and the retroactive energy mix cannot be known with total accuracy and realism;
 21 thus, simplifying yet tenable assumptions are necessary and justified.

22 To determine the proportional substitution by coal and gas in our baseline historical
 23 scenario, we first examine the world nuclear reactor properties provided by IAEA¹². Based on
 24 typical international values for coal and gas capacity factors (CFs)¹³, we then assume that each of
 25 the 441 reactors listed in Table 14 of ref 12 with CF > 65% is replaced by coal, and each reactor
 26 with CF ≤ 65% is replaced by gas.

27 For each country x we first calculate $P_i(x)$, the power (*not* energy) generated by each
 28 reactor i:

$$29 \quad P_i(x) = CF_i(x) \times C_i(x) \quad (1)$$

30 where CF_i , C_i denote the reactor capacity factor and net capacity, respectively, listed in Table 14
 31 of ref 12. We then calculate $f_i(x)$, the CF-weighted proportion of generated power by each
 32 reactor:

$$33 \quad f_i(x) = P_i(x) / \sum_i P_i(x) \quad (2)$$

34 Next, we calculate $F_j(x)$, the total proportion of generated nuclear power replaced by power from
 35 fossil fuel j:

$$36 \quad F_j(x) = \sum_i f_i^{(j)}(x) \quad (3)$$

37 where $f_i^{(j)}(x)$ simply denotes grouping of all the f_i values by replacement fuel j. For reference, on
 38 the global scale this yields about 95% replacement by coal and 5% by gas in our baseline
 39 historical scenario, which we suggest is plausible for the reasons given in the Discussion section.
 40 Lastly, we calculate $I(x,t)$, the annual net prevented impacts (mortality or GHG emissions) from
 41 nuclear power in country x and year t as follows:

$$42 \quad I(x,t) = \sum_j [IF_j \times F_j(x) \times n(x,t)] - IF_n \times n(x,t) \quad (4)$$

43 where IF_j = impact factor for fossil fuel j (from Table 1); $n(x,t)$ = nuclear power generation (in
 44 energy units); and IF_n = impact factor for nuclear power (from Table 1). Note that the first term
 45 in eq 4 reflects gross avoided impacts, while the second reflects direct impacts of nuclear power.

46 For the projection period (2010-2050), using eq 4 we calculate human deaths and GHG
 47 emissions that could result if all projected nuclear power production is canceled and again
 48 replaced only by fossil fuels. Of course, some or most of this hypothetically canceled nuclear
 49 power could be replaced by power from renewables, which have generally similar impact factors
 50 as nuclear (e.g., see Figure 2 of ref 7). Thus, our results for the projection period should
 51 ultimately be viewed as upper limits on potentially prevented impacts from future nuclear power.

1 We project annual nuclear power production in the regions containing the top five CO₂-
2 emitting countries and Western Europe based on the regional decadal projections in Table 4 of
3 IAEA⁶, which we linearly interpolate to an annual scale. To set $F_j(x)$ in eq 4, we consider two
4 simplified cases for both the global and regional scales. In the first (“All Coal”), $F_j(x)$ is fixed at
5 100% coal, and in the second (“All Gas”) it is fixed at 100% gas. This approach yields the full
6 range of potentially prevented impacts from future nuclear power. It is taken here because of the
7 lack of country-specific projections in ref 6 as well as the large uncertainty in determining which
8 fossil fuel(s) could replace *future* nuclear power, given recent trends in electricity production
9 (Figure 1, Figure S3, and ref 14).
10
11

12 **Methodological limitations**

14 The projections for nuclear power by IAEA⁶ assume essentially no climate change
15 mitigation measures in the low-end case and aggressive mitigation measures in the high-end
16 case. It is unclear which path the world will follow – however, these IAEA projections *do* take
17 into account the effects of the Fukushima accident. It seems that except possibly for Japan, the
18 top five CO₂-emitting countries are not planning a phase-down of pre-Fukushima plans for future
19 nuclear power. For instance China, India, and Russia have affirmed plans to increase their
20 current nuclear capacity by >3-fold, >12-fold, and 2-fold, respectively (see Table 12.2 of ref 2).
21 In Japan the future of nuclear power now seems unclear – as of ~1 year following the Fukushima
22 accident, nuclear power generation in Japan decreased by 63%, while fossil fuel power
23 generation increased by 26% (ref 15), thereby almost certainly increasing Japan’s CO₂
24 emissions.
25
26

27 Although our analysis reflects mortality from all stages of the fuel cycle for each energy
28 source, it excludes serious illnesses, including respiratory and cerebrovascular hospitalizations,
29 chronic bronchitis, congestive heart failure, non-fatal cancers, and hereditary effects. For fossil
30 fuels, such illnesses are estimated to be approximately 10 times higher than the mortality factors
31 in Table 1, while for nuclear power they are ~3 times higher¹⁶. Another important limitation is
32 that the mortality factors exclude the impacts of anthropogenic climate change and development-
33 related differences, as explained in the Discussion section. Aspects of nuclear power that cannot
34 meaningfully be quantified due to very large uncertainties (e.g. potential mortality from
35 proliferation of weapons-grade material) are also not included in our analysis.
36
37

38 Proportions of fossil fuels in our projection cases are assumed to be fixed (for the purpose
39 of determining upper and lower bounds), but will almost certainly vary across years and decades,
40 as in the historical period (Figure 1). The dominance of coal in the global average electricity mix
41 seems likely for the near future though (e.g., Figure 5.2 of ref 2). However, even if there is large-
42 scale worldwide electric fuel switching from coal to gas, our assessment is that the ultimate GHG
43 savings from such a transition are unlikely to be sufficient to minimize the risk of dangerous
44 anthropogenic climate change (unless the resulting emissions are captured and stored), as
45 discussed in the next section.
46
47
48

49 **Results and Discussion**

50 **Mortality**

53 We calculate a mean value of 1.84 million human deaths prevented by world nuclear
54 power production from 1971-2009 (see Figure 2a for full range), with an average of 76,000
55 prevented deaths/yr from 2000-2009 (range 19,000–300,000). Estimates for the top five CO₂
56 emitters, along with full estimate ranges for all regions in our baseline historical scenario, are
57 also shown in Figure 2a. For perspective, results for upper and lower bound scenarios are shown
58
59
60

1 in Figure S1. In Germany, which has announced plans to shut down all reactors by 2022 (ref 2),
2 we calculate that nuclear power has prevented an average of over 117,000 deaths from 1971-
3 2009 (range 29,000–470,000). The large ranges stem directly from the ranges given in Table 1
4 for the mortality factors.

5 Our estimated human deaths *caused* by nuclear power from 1971-2009 are far lower than
6 the avoided deaths. Globally, we calculate 4900 such deaths, or about 370 times lower than our
7 result for avoided deaths. Regionally, we calculate approximately 1,800 deaths in OECD Europe;
8 1,500 in USA; 540 in Japan; 460 in Russia (includes all 15 Former Soviet Union countries); 40
9 in China; and 20 in India. About 25% of these deaths are due to occupational accidents and about
10 70% are due to air pollution-related effects (presumably fatal cancers from radiation fallout; see
11 Table 2 of ref 16).

12 However, empirical evidence indicates that the April 1986 Chernobyl accident was the
13 world's only source of fatalities from nuclear power plant radiation fallout. According to the
14 latest assessment by the United Nations Scientific Committee on the Effects of Atomic Radiation
15 (UNSCEAR)¹⁷, 43 deaths are conclusively attributable to radiation from Chernobyl as of 2006
16 (28 were plant staff/first responders and 15 were from the 6,000 diagnosed cases of thyroid
17 cancer). UNSCEAR¹⁷ also states that reports of an increase in leukemia among recovery workers
18 who received higher doses are inconclusive, although cataract development was clinically
19 significant in that group; otherwise, for these workers as well as the general population, “there
20 has been no persuasive evidence of any other health effect” attributable to radiation exposure¹⁷.

21 Furthermore, no deaths have been conclusively attributed (in a scientifically valid manner)
22 to radiation from the other two major accidents, i.e. Three Mile Island in March 1979, for which
23 a 20-year comprehensive scientific health assessment was done¹⁸; and the March 2011
24 Fukushima Daiichi accident. While it is too soon to meaningfully assess the health impacts of the
25 latter accident, one early analysis¹⁹ indicates that annual radiation doses in nearby areas were
26 much lower than the generally accepted 100 mSv threshold¹⁷ for fatal disease development. In
27 any case, our calculated value for global deaths caused by historical nuclear power (~4,900)
28 could be a major overestimate relative to the empirical value (by 2 orders of magnitude). The
29 absence of evidence of large mortality from past nuclear accidents is consistent with recent
30 findings^{20, 21} that the “linear no-threshold” model used to derive the nuclear mortality factor in
31 Table 1 (see ref 22) might not be valid for the relatively low radiation doses that the public was
32 exposed to from nuclear power plant accidents.

33 For the projection period (2010-2050), we find that in the All Coal case (see Methods),
34 an average of 4.39 million and 7.04 million deaths are prevented globally by nuclear power
35 production for the low-end and high-end projections of IAEA⁶, respectively, while in the All Gas
36 case, an average of 420,000 and 680,000 deaths are prevented globally (see Figure 2b,c for full
37 ranges). Regional results are also shown in Figure 2b,c. The Far East and North America have
38 particularly high values, given that they are projected to be the biggest nuclear power producers
39 (Figure S2). As in the historical period, calculated deaths caused by nuclear power in our
40 projection cases are far lower (2 orders of magnitude) than the avoided deaths, even taking the
41 nuclear mortality factor at face value (despite the discrepancy with empirical data discussed
42 above for the historical period).

43 The substantially lower deaths in the projected All Gas case follow simply from the fact
44 that gas is estimated to have a mortality factor an order of magnitude lower than coal (Table 1).
45 However, this does not necessarily provide a valid argument for such large-scale “fuel
46 switching” for mitigation of either climate change or air pollution, for several reasons. First, it is
47 important to bear in mind that our results for prevented mortality are likely conservative, because
48 the mortality factors in Table 1 do not incorporate impacts of ongoing or future anthropogenic
49 climate change¹⁶. These impacts are likely to become devastating for both human health and
50

ecosystems if recent global GHG emission trends continue^{1,3}. Also, potential global natural gas resources are enormous – e.g., published estimates for technically recoverable unconventional gas resources suggest a carbon content ranging from >700 GtCO₂ (based on refs 23, 24) to >17,000 GtCO₂ (based on refs 24, 25). While we acknowledge that natural gas might play an important role as a “transition” fuel to a clean-energy era due to its lower mortality (and emission) factor relative to coal, we stress that long-term, widespread use of natural gas (without accompanying carbon capture and storage) could lead to unabated GHG emissions for many decades, given the typically multi-decadal lifetime of energy infrastructure -- thereby greatly complicating climate change mitigation efforts.

GHG emissions

We calculate that world nuclear power generation prevented an average of 64 GtCO₂-eq (or 17 GtC-eq) of cumulative emissions from 1971-2009 (Figure 3a; see full range therein), with an average of 2.6 GtCO₂-eq/yr prevented annual emissions from 2000-2009 (range 2.4–2.8 GtCO₂/yr). Regional results are also shown in Figure 3a. Our global results are 7-14% lower than previous estimates^{8,9} that, among other differences, assumed all historical nuclear power would have been replaced only by coal; and 34% higher than in another study¹⁰ in which the methodology is not explained clearly enough to infer the basis for the differences. Given that cumulative and annual global fossil fuel CO₂ emissions during the above periods were 840 GtCO₂ and 27 GtCO₂/yr respectively¹¹, our mean estimate for cumulative prevented emissions may not appear substantial – however it is instructive to look at other quantitative comparisons.

For instance, 64 GtCO₂-eq amounts to the cumulative CO₂ emissions from coal burning over approximately the past 35 yr in USA, 17 yr in China, or 7 yr in the top five CO₂ emitters¹⁷. Also, since a 500-MW coal-fired power plant typically emits 3 MtCO₂/yr²⁶, 64 GtCO₂-eq is equivalent to the cumulative lifetime emissions from almost 430 such plants, assuming an average plant lifetime of 50 yr. It is therefore evident that without global nuclear power generation in recent decades, near-term mitigation of anthropogenic climate change would pose a much greater challenge.

For the projection period (2010-2050), in the All Coal case an average of 150 and 240 GtCO₂-eq cumulative global emissions are prevented by nuclear power for the low-end and high-end projections of IAEA⁶, respectively, while in the All Gas case, an average of 80 and 130 GtCO₂-eq emissions are prevented (see Figure 3b,c for full ranges). Regional results are also shown in Figure 3b,c. These results also differ substantially from previous studies^{9,10}, largely due to differences in nuclear power projections (see SI).

To put our calculated overall mean estimate (80-240 GtCO₂-eq) of potentially prevented future emissions in perspective, note that to achieve a 350 ppm CO₂ target near the end of this century, cumulative “allowable” fossil CO₂ emissions from 2012 to 2050 are at most ~500 GtCO₂ (ref 3). Thus, projected nuclear power could reduce the climate change mitigation burden by 16-48% over the next few decades (derived by dividing 80 and 240 by 500).

Uncertainties

Our results contain various uncertainties, primarily stemming from our impact factors (Table 1) and our assumed replacement scenarios for nuclear power. In reality the impact factors are not likely to remain static, as we (implicitly) assumed – for instance, emission factors depend heavily on the particular mix of energy sources. Because the factors in Table 1 neglect ongoing or projected climate impacts as well as the significant disparity in pollution between developed and developing countries¹⁶, our results for both avoided GHG emissions and avoided mortality could be substantial underestimates. For example, in China, where coal burning accounts for over 75% of electricity generation in recent decades (ref 14; Figure S3), some coal-fired power

1 plants that meet domestic environmental standards have a mortality factor almost 3 times higher
2 than the mean global value (Table 1). These differences related to development status will
3 become increasingly important as fossil fuel use and GHG emissions of developing countries
4 continue to outpace those of developed countries¹¹.

5
6 On the other hand, if coal would not have been as dominant a replacement for nuclear as
7 assumed in our baseline historical scenario, then our avoided historical impacts could be
8 overestimates, since coal causes much larger impacts than gas (Table 1). However, there are
9 several reasons this is unlikely. Key characteristics of coal plants (e.g., plant capacity, capacity
10 factor, and total production costs) are historically much more similar to nuclear plants than are
11 those of natural gas plants¹³. Also, the vast majority of existing nuclear plants were built before
12 1990, but advanced gas plants that would be suitable replacements for base-load nuclear plants
13 (i.e., combined-cycle gas turbines) have only become available since the early 1990s¹³.
14 Furthermore, coal resources are highly abundant and widespread^{24,25} and coal fuel and total
15 production costs have long been relatively low, unlike historically available gas resources and
16 production costs¹³. Thus, it is not surprising that coal has been by far the dominant source of
17 global electricity thus far (Figure 1). We therefore assess that our baseline historical replacement
18 scenario is plausible and that it is not as significant an uncertainty source as the impact factors –
19 i.e., our avoided historical impacts are more likely underestimates, as discussed in the above
20 paragraph.
21
22
23
24

25 Implications

26 More broadly, our results underscore the importance of avoiding a false and
27 counterproductive dichotomy between reducing air pollution and stabilizing the climate, as
28 elaborated by others²⁷⁻²⁹. If near-term air pollution abatement trumps the goal of long-term
29 climate protection, governments might decide to carry out future electric fuel switching in much
30 more climate-impacting ways than we have examined here. For instance, they might start large-
31 scale production and use of gas derived from coal ("syngas"), as coal is by far the most abundant
32 of the three conventional fossil fuels^{24,25}. While this could reduce the very high pollution-related
33 deaths from coal power (Figure 2), the GHG emissions factor for syngas is substantially higher
34 (between ~5-90%) than for coal³⁰, thereby entailing even higher electricity sector GHG
35 emissions in the long term.
36
37

38 In conclusion, it is clear that nuclear power has provided a large contribution to the
39 reduction of global mortality and GHG emissions due to fossil fuel use. If the role of nuclear
40 power significantly declines in the next few decades, the International Energy Agency asserts
41 that achieving a target atmospheric GHG level of 450 ppm CO₂-eq would require "heroic
42 achievements in the deployment of emerging low-carbon technologies, which have yet to be
43 proven. Countries that rely heavily on nuclear power would find it particularly challenging and
44 significantly more costly to meet their targeted levels of emissions."² Our analysis herein and a
45 prior one⁷ strongly support this conclusion. Indeed, based on combined evidence from
46 paleoclimate data, observed ongoing climate impacts, and the measured planetary energy
47 imbalance, it appears increasingly clear that the commonly discussed targets of 450 ppm and 2°C
48 global temperature rise (above preindustrial levels) are insufficient to avoid devastating climate
49 impacts – we have suggested elsewhere that more appropriate targets are <350 ppm and 1°C
50 (refs 3, 31-33). Achieving these targets emphasizes the importance of retaining and expanding
51 nuclear power, as well as carbon-free renewables, in the near-term global energy supply.
52
53
54
55
56
57
58
59
60

References

1. Parry, M. L.; Canziani, O. F.; Palutikof, J. P.; van der Linden, J.; Hanson, C. E., Eds. *Climate Change 2007: Impacts, Adaptation and Vulnerability*; Contribution of Working Group III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change (IPCC); Cambridge Univ. Press: Cambridge, 2007.
2. *World Energy Outlook 2011*; OECD/International Energy Agency, 2011.
3. Hansen, J.; Kharecha, P.; Sato, M.; Ackerman, F.; Hearty, P. J.; Hoegh-Guldberg, O.; Hsu, S-L.; Krueger, F.; Parmesan, C.; Rahmstorf, S.; Rockstrom, J.; Rohling, E. J.; Sachs, J.; Smith, P.; Steffen, K.; Van Susteren, L.; von Schuckmann, K.; Zachos, J. C., *et al.*, Scientific case for avoiding dangerous climate change to protect young people and nature. *Proc. Natl Acad. Sci. USA* (submitted).
4. Global Health Observatory Data Repository; World Health Organization, 2012; <http://apps.who.int/gho/data/?vid=34300> (accessed November 2012)
5. *Climate Change and Nuclear Power 2011*; International Atomic Energy Agency, 2011; http://www.iaea.org/OurWork/ST/NE/Pess/assets/11-43751_ccnp_brochure.pdf (accessed September 2012)
6. *Energy, Electricity and Nuclear Power Estimates for the Period up to 2050: 2011 Edition*; International Atomic Energy Agency, 2011; http://www-pub.iaea.org/MTCD/Publications/PDF/RDS1_31.pdf (accessed January 2012)
7. Kharecha, P. A.; Kutscher, C. F.; Hansen, J. E.; Mazria, E. Options for near-term phaseout of CO₂ emissions from coal use in the United States. *Environ. Sci. Technol.* **2010**, *44*, 4050-4062.
8. *Nuclear Energy and Addressing Climate Change*; Organization for Economic Cooperation and Development, 2009; <http://www.oecd-nea.org/press/in-perspective/addressing-climate-change.pdf> (accessed February 2012)
9. Lenzen, M.; Schaeffer, R. Historical and potential future contributions of power technologies to global warming. *Clim. Ch.* **2011**, doi 10.1007/s10584-011-0270-y.
10. Coleman, N. M.; Abramson, L. R.; Coleman, F. A. B. Estimated lag time in global carbon emissions and CO₂ concentrations produced by commercial nuclear power through 2009 with projections through 2030. *Health Physics* **2012**, *102*, 326-334.
11. Boden, T. A.; Marland, G.; Andres, R. J. Global, Regional, and National Fossil-Fuel CO₂ Emissions. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge, Tenn., U.S.A., 2012, doi 10.3334/CDIAC/00001_V2012.
12. *Nuclear Power Reactors in the World: 2011 Edition*; International Atomic Energy Agency, 2011; <http://www-pub.iaea.org/books/iaeabooks/8752/Nuclear-Power-Reactors-in-the-World-2011-Edition> (accessed July 2012)
13. See individual technology briefs for coal, gas, and nuclear power by the International Energy Agency- Energy Technology Systems Analysis Program (IEA-ETSAP), available at <http://www.iea-etsap.org/web/E-techDS.asp> (accessed July 2012)
14. *IEA World Energy Statistics and Balances* (database); International Energy Agency, 2010; doi: 10.1787/data-00512-en.
15. Electricity Generated and Purchased in FY 2011; Federation of Electric Power Companies of Japan, 2012; http://www.fepec.or.jp/english/news/generated_purchased/_icsFiles/afieldfile/2012/04/19/hat_suju_E_FY2011.pdf (accessed September 2012)
16. Markandya, A.; Wilkinson, P. Electricity generation and health. *Lancet* **2007**, *370*, 979-990.
17. *Sources and Effects of Ionizing Radiation. Annex D: Health effects due to radiation from the Chernobyl accident*; United Nations Scientific Committee on the Effects of Atomic

- Radiation, 2011; http://www.unscear.org/docs/reports/2008/11-80076_Report_2008_Annex_D.pdf (accessed August 2011)
18. Talbott, E. O.; Youk, A. O.; McHugh-Pemu, K. P.; Zborowski, J. V. Long-term follow-up of the residents of the Three Mile Island Accident area: 1979-1998. *Environ. Health Perspect.* **2003**, *111*, 341-348.
 19. Kinoshita, N.; Sueki, K.; Sasa, K.; Kitagawa, J.; Ikarashi, S.; Nishimura, T.; Wong, Y-S.; Satou, Y.; Handa, K.; Takahashi, T.; Sato, M.; Yamagata, T. Assessment of individual radionuclide distributions from the Fukushima nuclear accident covering central-east Japan. *Proc. Natl. Acad. Sci. USA* **2011**, *108*, 19526-19529.
 20. Tanooka, H. Meta-analysis of non-tumour doses for radiation-induced cancer on the basis of dose-rate. *Int. J. Radiat. Biol.* **2011**, *87*, 645-652.
 21. Neumaier, T.; Swenson, J.; Pham, C.; Polyzos, A.; Lod, A. T.; Yang, P.; Dyball, J.; Asaithamby, A.; Chen, D. J.; Bissell, M. J.; Thalhammer, A.; Costes, S. V. Evidence for formation of DNA repair centers and dose-response nonlinearity in human cells. *Proc. Natl. Acad. Sci. USA* **2012**, *109*, 443-448.
 22. *ExternE: Externalities of Energy. Methodology 2005 Update*; European Communities, 2005; <http://www.externe.info/publications.html> (accessed February 2012)
 23. *Are We Entering a Golden Age of Gas? Special Report, World Energy Outlook 2011*; OECD/International Energy Agency, 2011; http://www.iea.org/weo/docs/weo2011/WEO2011_GoldenAgeofGasReport.pdf (accessed June 2011)
 24. Metz, B.; Davidson, O. R.; Bosch, P. R.; Dave, R.; Meyer, L. A., Eds. *Climate Change 2007: Mitigation*; Contribution of Working Group III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change (IPCC); Cambridge Univ. Press: Cambridge, 2007.
 25. *Global Energy Assessment – Toward a Sustainable Future*; International Institute for Applied Systems Analysis, 2012; <http://www.globalenergyassessment.org> (accessed June 2012)
 26. *The Future of Coal*; Massachusetts Institute of Technology, 2007; <http://web.mit.edu/coal> (accessed December 2007)
 27. Hansen, J.; Sato, M.; Ruedy, R.; Lacis, A.; Oinas, V. Global warming in the twenty-first century: An alternative scenario. *Proc. Natl. Acad. Sci. USA* **2000**, *97*, 9875-9880.
 28. Ramanathan, V.; Feng, Y. Air pollution, greenhouse gases and climate change: Global and regional perspectives. *Atmos. Environ.* **2009**, *43*, 37-50.
 29. Shindell, D.; Kuylenstierna, J.C.I.; Vignati, E.; van Dingenen, R.; Amann, M.; Klimont, Z.; Anenberg, S.C.; Muller, N.; Janssens-Maenhout, G.; Raes, F.; Schwartz, J.; Faluvegi, G.; Pozzoli, L.; Kupiainen, K.; Höglund-Isaksson, L.; Emberson, L.; Streets, D.; Ramanathan, V.; Hicks, K.; Oanh, N.T.K.; Milly, G.; Williams, M.; Demkine, V.; Fowler, D. Simultaneously mitigating near-term climate change and improving human health and food security. *Science* **2012**, *335*, 183-189.
 30. Jaramillo, P.; Griffin, W. M.; Matthews, H. S. Comparative life-cycle air emissions of coal, domestic natural gas, LNG, and SNG for electricity generation. *Environ. Sci. Technol.* **2007**, *41*, 6290-6296.
 31. Hansen, J.; Sato, M.; Kharecha, P.; Beerling, D.; Berner, R.; Masson-Delmotte, V.; Pagani, M.; Raymo, M.; Royer, D. L.; and Zachos, J. C. Target atmospheric CO₂: Where should humanity aim? *Open Atmos. Sci. J.* **2008**, *2*, 217-231.
 32. Rockstrom, J.; Steffen, W.; Noone, K.; Persson, A.; Chapin, F. S.; III, Lambin, E.; Lenton, T. M.; Scheffer, M.; Folke, C.; Schellnhuber, H.; Nykvist, B.; De Wit, C. A.; Hughes, T.; van der Leeuw, S.; Rodhe, H.; Sorlin, S.; Snyder, P. K.; Costanza, R.; Svedin, U.; Falkenmark,

- 1 M.; Karlberg, L.; Corell, R. W.; Fabry, V. J.; Hansen, J.; Walker, B.; Liverman, D.;
2 Richardson, K.; Crutzen, P.; and Foley, J. A safe operating space for humanity. *Nature* **2009**,
3 *461*, 472-475, doi:10.1038/461472a.
4
5 33. Hansen, J.; Sato, M.; Kharecha, P.; von Schuckmann, K. Earth's energy imbalance and
6 implications. *Atmos. Chem. Phys.* **2011**, *11*, 13421-13449 doi:10.5194/acp-11-13421-2011.
7
8 34. Lenzen, M. Life cycle energy and greenhouse gas emissions of nuclear energy: A review.
9 *Energ. Convers. Manage.* **2008**, *49*, 2178-2199.
10
11 35. Lenzen, M. Current State of Development of Electricity-Generating Technologies: A
12 Literature Review. *Energies* **2010**, *3*, 462-591.
13

14 **Acknowledgements:**

15 We thank Chuck Kutscher of the U.S. DOE-National Renewable Energy Laboratory for helpful
16 comments on our methodology and three anonymous reviewers for helpful feedback on our
17 manuscript. Funding for this work was provided by the Lenfest Foundation and the Columbia
18 University-NASA Cooperative Agreement (Award NNX11AR63A). P.K. designed the study
19 with input from J.H.; P.K. performed the calculations and analysis, and wrote the paper with
20 feedback from J.H.
21
22
23
24

25 **Supporting Information Available**

26 Supplemental discussion and three supplemental figures accompany this article. This information
27 is available free of charge via the Internet at <http://pubs.acs.org/>.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1. World electricity generation by power source for 1971-2009 (data from ref 14). In the last decade (2000-2009), nuclear power provided an average 15% of world generation; coal, gas, and oil provided 40%, 20%, and 6%, respectively; and renewables provided 16% (hydropower) and 2% (non-hydro).




Figure 2. Cumulative net deaths prevented assuming nuclear power replaces fossil fuels. Results for (a) the historical period in our study (1971-2009), showing mean values (labeled) and ranges for the baseline historical scenario. Results for (b) the high-end and (c) low-end projections of nuclear power production by the UN International Atomic Energy Agency⁶ for the period 2010-2050. Error bars reflect the ranges for the fossil fuel mortality factors listed in Table 1. The larger columns in panels b and c reflect the All Coal case and are labeled with their mean values, while the smaller columns reflect the All Gas case; values for the latter are not shown because they are all simply a factor of ~ 10 lower (reflecting the order-of-magnitude difference between the mortality factors for coal and gas shown in Table 1). Countries/regions are arranged in descending order of CO₂ emissions in recent years. FSU15=15 countries of the Former Soviet Union and OECD=Organization for Economic Cooperation and Development.




Figure 3. Cumulative net GHG emissions prevented assuming nuclear power replaces fossil fuels. Same panel arrangement as Figure 2, except mean values for all cases are labeled. Error bars reflect the ranges for the fossil fuel emission factors listed in Table 1.

Table 1. Mortality and GHG emission factors used in this study. Mortality factors are based on analysis for Europe (except as indicated) and represent the sum of accidental deaths and air pollution-related effects in Table 2 of ref 16. They reflect impacts from all stages of the fuel cycle, including fuel extraction, transport, transformation, waste disposal, and electricity transport. Their ranges are 95% confidence intervals and represent deviation from the mean by a factor of ~4. Mortality factor for coal is the mean of the factors for lignite and coal in ref 16. Mean values for emission factors are the midpoints of the ranges given in the sources. Water pollution is also a significant impact but is not factored into these values. Additional uncertainties and limitations inherent in these factors are discussed in the text.

Electricity source	Mean value (<i>Range</i>)	Unit	Source
Coal	28.67 (7.15-114)	deaths/TWh	ref 16
	77 (19.25-308)	deaths/TWh	ref 16 (China)*
	1045 (909-1182)	tCO ₂ -eq/GWh	ref 30
Natural gas	2.821 (0.7-11.2)	deaths/TWh	ref 16
	602 (386-818)	tCO ₂ -eq/GWh	ref 30
Nuclear	0.074 (<i>range not given</i>)	deaths/TWh	ref 16
	65 (10-130)**	tCO ₂ -eq/GWh	ref 34

TWh = terawatt-hr, GWh = gigawatt-hr; tCO₂-eq = tonnes CO₂-equivalent emissions.

*Range not given in source for China, but for consistency with other factors, assumed to be 4 times lower and higher than mean

**Some authors contend the upper limit is significantly higher, but their conclusions are based on dubious assumptions³⁵




Figure 1. World electricity generation by power source for 1971-2009 (data from ref 14). In the last decade (2000-2009), nuclear power provided an average 15% of world generation; coal, gas, and oil provided 40%, 20%, and 6%, respectively; and renewables provided 16% (hydropower) and 2% (non-hydro).
127x82mm (300 x 300 DPI)




Figure 2. Cumulative net deaths prevented assuming nuclear power replaces fossil fuels. Results for (a) the historical period in our study (1971-2009), showing mean values (labeled) and ranges for the baseline historical scenario. Results for (b) the high-end and (c) low-end projections of nuclear power production by the UN International Atomic Energy Agency⁶ for the period 2010-2050. Error bars reflect the ranges for the fossil fuel mortality factors listed in Table 1. The larger columns in panels b and c reflect the All Coal case and are labeled with their mean values, while the smaller columns reflect the All Gas case; values for the latter are not shown because they are all simply a factor of ~ 10 lower (reflecting the order-of-magnitude difference between the mortality factors for coal and gas shown in Table 1). Countries/regions are arranged in descending order of CO₂ emissions in recent years. FSU15=15 countries of the Former Soviet Union and OECD=Organization for Economic Cooperation and Development.

127x82mm (300 x 300 DPI)




Figure 3. Cumulative net GHG emissions prevented assuming nuclear power replaces fossil fuels. Same panel arrangement as Figure 2, except mean values for all cases are labeled. Error bars reflect the ranges for the fossil fuel emission factors listed in Table 1.
127x82mm (300 x 300 DPI)

Number of deaths prevented annually by nuclear power
1971-2009

